PPE COVID-19 Guidance Updated 5/1/2020						
PROCEDURES	ANESTHESIA	ST	CIRCULATOR	PRE-OP	POST-OP STAFF	OR AIR FLOW/CLEAN UP
ENT - BMV	Eye shield N95 mask New gown	Eye shield N95 mask New gown	Eye shield N95 mask New gown	Surgical mask	Phase I Recovery Eye shield N95 mask Phase II Surgical mask	<ul> <li>Step 1 Remove anesthesia circuit from room</li> <li>Step 2 Remove contaminated instruments</li> <li>Step 3 Wait 15 minutes</li> <li>Step 4 Clean room and set up</li> </ul>
ENT - ALL OTHER CASES	Eye shield N95 mask New gown	Eye shield N95 mask New gown	Eye shield N95 mask New gown	Surgical mask	Phase I Recovery Eye shield N95 mask Phase II Surgical mask	<ul> <li>Step 1 Remove anesthesia circuit from room</li> <li>Step 2 Remove contaminated instruments</li> <li>Step 3 Wait 15 minutes</li> <li>Step 4 Clean room and set up</li> </ul>
ORTHO/PLASTICS General/MAC/Spinal	Eye shield N95 mask New gown	Eye shield N95 mask New gown	Eye shield N95 mask New gown	Surgical mask	Phase I Recovery Eye shield N95 mask Phase II Surgical mask	<ul> <li>Step 1 Remove anesthesia circuit from room</li> <li>Step 2 Remove contaminated instruments</li> <li>Step 3 Wait 15 minutes</li> <li>Step 4 Clean room and set up</li> </ul>
LOCAL/MR/IV Sedation	N/A	Eye shield surgical mask	Eye shield surgical mask	Surgical mask	Surgical mask	Normal turn over

<sup>\*</sup>NO gown for ST if procedure allows you to remain 6 feet away from patient throughout procedure.

<sup>\*</sup>ALL N95 users are to cover N95 with regular surgical mask and wear for 5 days.